


AHPWA22AUSA

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1(Original). A method of treating acne and/or hirsutism comprising the step of delivering to a mammal in need thereof a composition comprising a compound of formula I, or a tautomer thereof, and a physiologically compatible carrier, wherein formula I is:



wherein:

R^1 and R^2 are selected from the group consisting of H, alkyl, substituted alkyl, OH, O(alkyl), O(substituted alkyl), O(Acetyl), aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, alkylaryl, substituted alkylaryl, alkylheteroaryl, substituted alkylheteroaryl, 1-propynyl, substituted 1-propynyl, 3-propynyl, and substituted 3-propynyl;

or R^1 and R^2 are joined to form a ring selected from the group consisting of $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2-$, $-\text{O}(\text{CH}_2)_m\text{CH}_2-$, $-\text{O}(\text{CH}_2)_p\text{O}-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{N}(\text{H})\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{N}(\text{alkyl})\text{CH}_2\text{CH}_2-$;

m is an integer from 1 to 4;

n is an integer from 1 to 5;

p is an integer from 1 to 4;

or R^1 and R^2 form a double bond to $\text{C}(\text{CH}_3)_2$, $\text{C}(\text{cycloalkyl})$, O, or $\text{C}(\text{cycloether})$;

AHPWA22AUSA


R^3 is selected from the group consisting of H, OH, NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_6 alkenyl, substituted C_3 to C_6 alkenyl, alkynyl, substituted alkynyl, and COR^A ;

R^A is selected from the group consisting of H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, and substituted C_1 to C_3 aminoalkyl;

R^4 is selected from the group consisting of H, halogen, CN, NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_1 to C_6 alkoxy, substituted C_1 to C_6 alkoxy, C_1 to C_6 aminoalkyl, and substituted C_1 to C_6 aminoalkyl;

R^5 is selected from the group consisting of a), b) and c):

a) a substituted benzene ring having the structure:



X is selected from the group consisting of halogen, OH, CN, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 thioalkyl, substituted C_1 to C_3 thioalkyl, $S(O)alkyl$, $S(O)_2alkyl$, C_1 to C_3 aminoalkyl, substituted C_1 to C_3 aminoalkyl, NO_2 , C_1 to C_3 perfluoroalkyl, substituted C_1 to C_3 perfluoroalkyl, 5 or 6 membered heterocyclic ring having 1 to 3 heteroatoms, $CONH_2$, $CSNH_2$, $CNHNHOH$, CNH_2NOH , $CNHNHOH$, COR^B , CSR^B , $OCOR^B$, and $NR^C COR^B$;

R^B is selected from the group consisting of H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, aryl, substituted aryl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, and substituted C_1 to C_3 aminoalkyl;

R^C is H, C_1 to C_3 alkyl, or substituted C_1 to C_3 alkyl;

Y and Z are independently selected from the group consisting of H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_4 alkyl, substituted C_1 to C_4 alkyl, C_1 to C_3 thioalkyl, and substituted C_1 to C_3 thioalkyl;

b) a five or six membered heterocyclic ring comprising 1, 2, or 3 heteroatoms selected from the group consisting of O, S, SO, SO_2 and NR^6 and having one or two

AHPWA22AUSA

independent substituents from the group consisting of H, halogen, CN, NO₂, C₁ to C₄ alkyl, substituted C₁ to C₄ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, COR^D, CSR^D, and NR^ECOR^D;

R^D is H, NH₂, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^E is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

R^F is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, or C₁ to C₄CO₂alkyl; or

c) an indol-4-yl, indol-7-yl or benzo-2-thiophene moiety, wherein said moiety is optionally substituted by from 1 to 3 substituents selected from the group consisting of halogen, alkyl, substituted alkyl, CN, NO₂, alkoxy, substituted alkoxy, and CF₃;


Qⁱ is S, NR⁷, or CR⁸R⁹;

R⁷ is selected from the group consisting of CN, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, SO₂CF₃, OR¹¹, and NR¹¹R¹²;

R⁸ and R⁹ are independent substituents selected from the group consisting of H, alkyl, substituted alkyl, acyl, substituted acyl, aroyl, substituted aroyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, NO₂, CN, and CO₂R¹⁰;

R¹⁰ is C₁ to C₃ alkyl or substituted C₁ to C₃ alkyl;

or CR⁸R⁹ comprise a six membered ring having the structure:



AHPWA22AUSA

R^{11} and R^{12} are independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, sulfonyl, and substituted sulfonyl; or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.


2(Original). The method according to Claim 1, wherein:

R^1 and R^2 are joined to form a $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$ ring;

n is 3;

R^3 and R^4 are H;

R^5 is the substituted benzene ring having the structure:



X is selected from the group consisting of halogen, CN, CONH_2 , CSNH_2 , COR^B , CSR^B , C_1 to C_3 alkoxy, C_1 to C_3 alkyl, NO_2 , C_1 to C_3 perfluoroalkyl, 5 membered heterocyclic ring comprising 1 to 3 heteroatoms, and C_1 to C_3 thioalkyl;

R^B is C_1 to C_3 aminoalkyl or substituted C_1 to C_3 aminoalkyl, wherein said aminoalkyl is $\text{NH}(\text{alkyl})$ or $\text{N}(\text{alkyl})_2$;

Y is selected from the group consisting of H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, C_1 to C_4 alkyl, and C_1 to C_3 thioalkyl.

3(Original). The method according to Claim 1, wherein:

R^1 and R^2 are joined to form the $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$ ring;

n is 3;

R^3 and R^4 are H;

AHPWA22AUSA

R^5 is the five membered ring having the structure:



U is O, S, or NR^6 ;

X' is selected from the group consisting of halogen, CN, NO_2 , $CONH_2$, $CSNH_2$, COR^B , CSR^B , C_1 to C_3 alkyl, and C_1 to C_3 alkoxy;

R^B is C_1 to C_3 aminoalkyl or substituted C_1 to C_3 aminoalkyl, wherein said aminoalkyl is $NH(alkyl)$ or $N(alkyl)_2$;

Y' is selected from the group consisting of H, halogen, and C_1 to C_4 alkyl, wherein said halogen is F.


4(Original). The method according to Claim 1, wherein:

R^1 and R^2 are joined to form a $-CH_2(CH_2)_nCH_2-$ ring;

n is 3;

R^3 and R^4 are H;

R^5 is the six membered ring having the structure:



X^1 is N or CX^2 ;

X^2 is halogen, CN, $CONH_2$, $CSNH_2$, COR^B , CSR^B , or NO_2 ;

R^B is C_1 to C_3 aminoalkyl or substituted C_1 to C_3 aminoalkyl, wherein said aminoalkyl is $NH(alkyl)$ or $N(alkyl)_2$.

5(Original). The method according to claim 1, wherein:

R^1 and R^2 are alkyl or substituted alkyl;

R^3 is H.

AHPWA22AUSA

6(Original). The method according to claim 1, wherein:

R^1 and R^2 are joined to form a ring selected from the group consisting of $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2-$, $-\text{O}(\text{CH}_2)_m\text{CH}_2-$, $-\text{O}(\text{CH}_2)_p\text{O}-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{N}(\text{H})\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{N}(\text{alkyl})\text{CH}_2\text{CH}_2-$;
 R^3 is H.

7(Original). The method according to claim 1, wherein:

R^3 is H;

Q^1 is S or NR^7 .

8(Original). The method according to claim 1, wherein the compound is delivered orally.

9(Currently Amended). The method according to ~~claims~~ claim 1, wherein said compound of formula I is selected from the group consisting of 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-thione, 3-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzonitrile, 4-1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-2-thiophenecarbonitrile, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile, 4-Methyl-5-(1,2-dihydro-2-thioxospiro[cyclohexane-1,3-[3H]-indol]-5-yl)-2-thiophenethioamide, 5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-(tert-butoxycarbonyl)-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-H-pyrrole-2-carbonitrile, 5-(2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methylpyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-3thiophenecarbonitrile, 5-(1,2-Dihydro-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-2thiophenecarbonitrile, 5-(3-Fluoro-4-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(2-Amino-5-pyrimidinyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-5-

AHPWA22AUSA

fluorobenzonitrile, 5-(3-chlorophenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-Benzyl-5-(3-chlorophenyl)-3-methyl-1,3-dihydro-2H-indole-2-thione, 4-(3,3-dimethyl-2-thioxo-2,3-dihydro-1H-indol-5-yl)-2-furonitrile, 5-(3-methoxyphenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-pyridinecarbonitrile, 5-(3,4-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(5-Chloro-2-thienyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-furancarbonitrile, 5-(3-Chloro-4-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chloro-5-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3,5-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-propyl-2-thiophenecarbonitrile, 5-(3-Fluoro-4-nitrophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-furancarbonitrile, 5''-(3-Chlorophenyl)spiro[cyclobutane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(2-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(4-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-4-methyl-2-thiophenecarbonitrile, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-2-thiophenecarbonitrile, 5''-(3-Fluorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5-(3-Hydroxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-chlorophenyl)-3,3-diethyl-1,3-dihydro-2H-indole-2-thione, 5-(4-Fluoro-3-(trifluoromethyl)phenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-n-butyl-2-thiophenecarbonitrile, 5-(3-Fluoro-5-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chlorophenyl)-N-hydroxyspiro[cyclohexane-1,3'-[3H]indol]-2-amine, N-(Acetyloxy)-5'-(3-chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2''amine, 5'-(3-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(2-

AHPWA22AUSA


Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(4-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3,4-difluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-methoxyphenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-nitrophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-cyanophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 3-(1',2'-Dihydro-2'-(hydroxyimino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluorobenzonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-1-methyl-2-carbonitrile, 5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-2-carbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'(acetoxymino)-5'-yl)-2-thiophenecarbonitrile, 3-Fluoro-N'-hydroxy-5-(2'-(hydroxyamino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarboximidamide, N'-Hydroxy-4-(spiro[cyclohexane-1,3'-[3H]indol]-2'-hydroxyimino)-5'-yl-2-thiophenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarboximidamide, 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(3-Cyano-5-fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-3-methyl-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-thiophen-3-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 3-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluoro-benzonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-1H-pyrrole-2-

AHPWA22AUSA

carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-4-methyl-thiophene-2-carbonitrile, and 4-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

10(Currently Amended). The method according to ~~claims~~ claim 1, wherein said compound is 5'-(5-Cyano-1-methyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

11(Original). A method of treating acne and/or hirsutism comprising the step of delivering to a mammal in need thereof a composition comprising a compound of formula II, or a tautomer thereof, and a physiologically compatible carrier, wherein formula II is:




wherein:

R¹¹ is selected from the group consisting of H, acyl, substituted acyl, aroyl, substituted aroyl, sulfonyl, and substituted sulfonyl;

R⁵ is (i), (ii), or (iii):

(i) a substituted benzene ring having the structure:



AHPWA22AUSA

wherein:

X is selected from the group consisting of halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, CNHNHOH, CNH₂NOH, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring comprising 1 to 3 heteroatoms, and C₁ to C₃ thioalkyl;

Y is selected from the group consisting of H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, and C₁ to C₃ thioalkyl;

(ii) a five membered ring having the structure:



wherein:


U is O, S, or NR⁶;

R⁶ is H, C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;

X' is selected from the group consisting of halogen, CN, NO₂, CONH₂, CNHNHOH, CNH₂NOH, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, C₁ to C₃ alkyl, and C₁ to C₃ alkoxy;

Y' is selected from the group consisting of H, F, and C₁ to C₄ alkyl; or

(iii) a six membered ring having the structure:



wherein:

X¹ is N or CX²;


X² is halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂ or NO₂;

or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

AHPWA22AUSA

12(Original). The method according to claim 11, wherein R^5 is said five membered ring and U is O or S.


13(Original). A method of treating acne and/or hirsutism comprising the step of delivering to a mammal in need thereof a composition comprising a compound of formula III, or a tautomer thereof, and a physiologically compatible carrier, wherein formula III is:



wherein:

R^5 is (i), (ii), or (iii):

(i) a substituted benzene ring having the structure:




wherein:

X is selected from the group consisting of halogen, CN, CONH_2 , CSNH_2 , CONHalkyl , CSNHalkyl , CON(alkyl)_2 , CSN(alkyl)_2 , CNHNOH , C_1 to C_3 alkoxy, C_1 to C_3 alkyl, NO_2 , C_1 to C_3 perfluoroalkyl, 5 membered heterocyclic ring comprising 1 to 3 heteroatoms, and C_1 to C_3 thioalkyl;

Y is selected from the group consisting of H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, C_1 to C_4 alkyl, and C_1 to C_3 thioalkyl;

(ii) a five membered ring having the structure:

AHPWA22AUSA



wherein:


U is O, S, or NR⁶;

R⁶ is H, C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;

X' is selected from the group consisting of halogen, CN, NO₂, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, C₁ to C₃ alkyl, and C₁ to C₃ alkoxy;

Y' is selected from the group consisting of H, F and C₁ to C₄ alkyl; or

(iii) a six membered ring having the structure:



wherein:

X¹ is N or CX²;


X² is halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂ or NO₂;

or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

14(Original). The method according to claim 13, wherein R⁵ is the five membered ring (ii) and U is O or S.

15(Original). A method of treating acne and/or hirsutism comprising the step of delivering to a mammal in need thereof a composition comprising a compound of formula IV, or a tautomer thereof, and a physiologically compatible carrier, wherein formula IV is:

AHPWA22AUSA




wherein:

R^8 is selected from the group consisting of H, CO_2R^{10} , acyl, substituted acyl, aroyl, substituted aroyl, alkyl, substituted alkyl, and CN;

R^{10} is C_1 to C_3 alkyl;

R^5 is (i), (ii), or (iii):

(i) a substituted benzene ring having the structure:




wherein:

X is selected from the group consisting of halogen, CN, CONH_2 , CSNH_2 , CONHalkyl , CSNHalkyl , CON(alkyl)_2 , CSN(alkyl)_2 , CNHNOH , C_1 to C_3 alkoxy, C_1 to C_3 alkyl, NO_2 , C_1 to C_3 perfluoroalkyl, 5 membered heterocyclic ring comprising 1 to 3 heteroatoms, and C_1 to C_3 thioalkyl;

Y is selected from the group consisting of H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, C_1 to C_4 alkyl, and C_1 to C_3 thioalkyl;

(ii) a five membered ring having the structure:



wherein:

U is O, S, or NR^6 ;


R^6 is H, C_1 to C_3 alkyl, or C_1 to C_4 CO_2alkyl ;

AHPWA22AUSA

X' is selected from the group consisting of halogen, CN, NO₂, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, C₁ to C₃ alkyl, and C₁ to C₃ alkoxy;

Y' is selected from the group consisting of H, F and C₁ to C₄ alkyl;

(iii) a six membered ring having the structure:



wherein:


X¹ is N or CX²;

X² is halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂ or NO₂;

or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

16(Original). The method according to claim 15, wherein R⁵ is the five-membered ring (ii) and U is O or S.


17(Original). A method of treating acne and hirsutism comprising the step of delivering to a mammal in need thereof a composition comprising a compound of formula V, or a tautomer thereof, and a physiologically compatible carrier, wherein formula V is:



R⁵ is (i), (ii), or (iii):

(i) a substituted benzene ring having the structure:

AHPWA22AUSA



wherein:

X is selected from the group consisting of halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, CNHNOH, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring comprising 1 to 3 heteroatoms, and C₁ to C₃ thioalkyl;

Y is selected from the group consisting of H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, and C₁ to C₃ thioalkyl;

(ii) a five membered ring having the structure:



wherein:


U is O, S, or NR⁶;

R⁶ is H, C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;

X' is selected from the group consisting of halogen, CN, NO₂, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, C₁ to C₃ alkyl, and C₁ to C₃ alkoxy;

Y' is selected from the group consisting of H, F, and C₁ to C₄ alkyl;

(iii) a six membered ring having the structure:



wherein:

X¹ is N or CX²;


AHPWA22AUSA

X^2 is halogen, CN, CONH_2 , CSNH_2 , CONHalkyl , CSNHalkyl , CON(alkyl)_2 , CSN(alkyl)_2 or NO_2 ;
or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

18(Original). The method according to claim 17, wherein R^5 is the five membered ring (ii) and U is O or S.

19(Withdrawn). A composition for conditioning the skin of a mammal in need thereof comprising:

- (i) a skin conditioning component; and
- (ii) a compound of formula I, or a tautomer thereof:



I

wherein:

R^1 and R^2 are selected from the group consisting of H, alkyl, substituted alkyl, OH, O(alkyl), O(substituted alkyl), O(Acetyl), aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, alkylaryl, substituted alkylaryl, alkylheteroaryl, substituted alkylheteroaryl, 1-propynyl, substituted 1-propynyl, 3-propynyl, and substituted 3-propynyl;

or R^1 and R^2 are joined to form a ring selected from the group consisting of $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2-$, $-\text{O}(\text{CH}_2)_m\text{CH}_2-$, $-\text{O}(\text{CH}_2)_p\text{O}-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{N}(\text{H})\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{N}(\text{alkyl})\text{CH}_2\text{CH}_2-$;

m is an integer from 1 to 4;

n is an integer from 1 to 5;

p is an integer from 1 to 4;

or R^1 and R^2 form a double bond to $\text{C}(\text{CH}_3)_2$, C(cycloalkyl), O, or C(cycloether);

AHPWA22AUSA


R^3 is selected from the group consisting of H, OH, NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_6 alkenyl, substituted C_3 to C_6 alkenyl, alkynyl, substituted alkynyl, and COR^A ;

R^A is selected from the group consisting of H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, and substituted C_1 to C_3 aminoalkyl;

R^4 is selected from the group consisting of H, halogen, CN, NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_1 to C_6 alkoxy, substituted C_1 to C_6 alkoxy, C_1 to C_6 aminoalkyl, and substituted C_1 to C_6 aminoalkyl;

R^5 is selected from the group consisting of a), b) and c):

a) a substituted benzene ring having the structure:



X is selected from the group consisting of halogen, OH, CN, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 thioalkyl, substituted C_1 to C_3 thioalkyl, $S(O)alkyl$, $S(O)_2alkyl$, C_1 to C_3 aminoalkyl, substituted C_1 to C_3 aminoalkyl, NO_2 , C_1 to C_3 perfluoroalkyl, substituted C_1 to C_3 perfluoroalkyl, 5 or 6 membered heterocyclic ring comprising 1 to 3 heteroatoms, $CONH_2$, $CSNH_2$, $CNHNHOH$, CNH_2NOH , $CNHNHOH$, COR^B , CSR^B , $OCOR^B$, and $NR^C COR^B$;

R^B is selected from the group consisting of H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, aryl, substituted aryl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, and substituted C_1 to C_3 aminoalkyl;

R^C is H, C_1 to C_3 alkyl, or substituted C_1 to C_3 alkyl;

Y and Z are independently selected from the group consisting of H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_4 alkyl, substituted C_1 to C_4 alkyl, C_1 to C_3 thioalkyl, and substituted C_1 to C_3 thioalkyl;

AHPWA22AUSA

b) a five or six membered heterocyclic ring comprising 1, 2, or 3 heteroatoms selected from the group consisting of O, S, SO, SO₂ and NR⁶ and having one or two independent substituents from the group consisting of H, halogen, CN, NO₂, C₁ to C₄ alkyl, substituted C₁ to C₄ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, COR^D, CSR^D, and NR^ECOR^D;

R^D is H, NH₂, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^E is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

R⁶ is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, or C₁ to C₄CO₂alkyl; or

c) an indol-4-yl, indol-7-yl or benzo-2-thiophene moiety, wherein said moiety is optionally substituted by from 1 to 3 substituents selected from the group consisting of halogen, alkyl, substituted alkyl, CN, NO₂, alkoxy, substituted alkoxy, and CF₃;


Q¹ is S, NR⁷, or CR⁸R⁹;

R⁷ is selected from the group consisting of CN, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, SO₂CF₃, OR¹¹, and NR¹¹R¹²;

R⁸ and R⁹ are independent substituents selected from the group consisting of H, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, NO₂, CN, and CO₂R¹⁰;

R¹⁰ is C₁ to C₃ alkyl or substituted C₁ to C₃ alkyl;

or CR⁸R⁹ comprise a six membered ring having the structure:



AHPWA22AUSA

R^{11} and R^{12} are independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, sulfonyl, and substituted sulfonyl; or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

20(Currently Amended). The ~~composition~~ method according to claim 19 22, wherein:

R^1 and R^2 are alkyl or substituted alkyl;
 R^3 is H.

21(Currently Amended). The ~~composition~~ method according to claim 19 22, wherein:

R^1 and R^2 are joined to form a ring selected from the group consisting of
-CH₂(CH₂)_nCH₂-, -CH₂CH₂C(CH₃)₂CH₂CH₂-, -O(CH₂)_mCH₂-, -O(CH₂)_pO-,
-CH₂CH₂OCH₂CH₂-, -CH₂CH₂N(H)CH₂CH₂-, and -CH₂CH₂N(alkyl)CH₂CH₂-;
 R^3 is H.


22(Currently Amended). The ~~composition~~ method according to claim 19 22, wherein:

R^3 is H;
 Q^1 is S or NR⁷.

23(Currently Amended). A method of conditioning the skin comprising the step of delivering to a mammal in need thereof a composition ~~according to claim~~ comprising:

- (i) a skin conditioning component; and
- (ii) a compound of formula I, or a tautomer thereof;

AHPWA22AUSA



wherein:

R¹ and R² are selected from the group consisting of H, alkyl, substituted alkyl, OH, O(alkyl), O(substituted alkyl), O(Acetyl), aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, alkylaryl, substituted alkylaryl, alkylheteroaryl, substituted alkylheteroaryl, 1-propynyl, substituted 1-propynyl, 3-propynyl, and substituted 3-propynyl;

or R¹ and R² are joined to form a ring selected from the group consisting of -CH₂(CH₂)_nCH₂-, -CH₂CH₂C(CH₃)₂CH₂CH₂-, -O(CH₂)_mCH₂-, -O(CH₂)_pO-, -CH₂CH₂OCH₂CH₂-, -CH₂CH₂N(H)CH₂CH₂-, and -CH₂CH₂N(alkyl)CH₂CH₂-;

m is an integer from 1 to 4;

n is an integer from 1 to 5;

p is an integer from 1 to 4;

or R¹ and R² form a double bond to C(CH₃)₂, C(cycloalkyl), O, or C(cycloether);

R³ is selected from the group consisting of H, OH, NH₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₆ alkenyl, substituted C₃ to C₆ alkenyl, alkynyl, substituted alkynyl, and COR^A;


R^A is selected from the group consisting of H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, and substituted C₁ to C₃ aminoalkyl;

R⁴ is selected from the group consisting of H, halogen, CN, NH₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₁ to C₆ alkoxy, substituted C₁ to C₆ alkoxy, C₁ to C₆ aminoalkyl, and substituted C₁ to C₆ aminoalkyl;

R⁵ is selected from the group consisting of a), b) and c):

a) a substituted benzene ring having the structure:

AHPWA22AUSA



X is selected from the group consisting of halogen, OH, CN, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ thioalkyl, substituted C₁ to C₃ thioalkyl, S(O)alkyl, S(O)₂alkyl, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, NO₂, C₁ to C₃ perfluoroalkyl, substituted C₁ to C₃ perfluoroalkyl, 5 or 6 membered heterocyclic ring comprising 1 to 3 heteroatoms, CONH₂, CSNH₂, CNHNHOH, CNH₂NOH, CNHNOH, COR^B, CSR^B, OCOR^B, and NR^CCOR^B.

R^B is selected from the group consisting of H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, and substituted C₁ to C₃ aminoalkyl;

R^C is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

Y and Z are independently selected from the group consisting of H, halogen, CN, NO₂, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₄ alkyl, substituted C₁ to C₄ alkyl, C₁ to C₃ thioalkyl, and substituted C₁ to C₃ thioalkyl;

b) a five or six membered heterocyclic ring comprising 1, 2, or 3 heteroatoms selected from the group consisting of O, S, SO, SO₂ and NR⁶ and having one or two independent substituents from the group consisting of H, halogen, CN, NO₂, C₁ to C₄ alkyl, substituted C₁ to C₄ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, COR^D, CSR^D, and NR^ECOR^D;

R^D is H, NH₂, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^E is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

R⁶ is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, or C₁ to C₄CO₂alkyl; or

c) an indol-4-yl, indol-7-yl or benzo-2-thiophene moiety, wherein said moiety is optionally substituted by from 1 to 3 substituents selected from the group

AHPWA22AUSA

consisting of halogen, alkyl, substituted alkyl, CN, NO₂, alkoxy, substituted alkoxy, and CF₃;


Q¹ is S, NR⁷, or CR⁸R⁹;

R⁷ is selected from the group consisting of CN, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, SO₂CF₃, OR¹¹, and NR¹¹R¹²;

R⁸ and R⁹ are independent substituents selected from the group consisting of H, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, NO₂, CN, and CO₂R¹⁰;

R¹⁰ is C₁ to C₃ alkyl or substituted C₁ to C₃ alkyl;

or CR⁸R⁹ comprise a six membered ring having the structure:



R¹¹ and R¹² are independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, sulfonyl, and substituted sulfonyl; or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

24(Currently Amended). The method according to ~~claims~~ claim 23 wherein said compound of formula I is selected from the group consisting of 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-thione, 3-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzonitrile, 4-1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-2-thiophenecarbonitrile, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile, 4-Methyl-5-(1,2-dihydro-2-thioxospiro[cyclohexane-1,3-[3H]-indol]-5-yl)-2-thiophenethioamide, 5-(1,2-

AHPWA22AUSA

Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-(tert-butoxycarbonyl)-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-H-pyrrole-2-carbonitrile, 5-(2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-3thiophenecarbonitrile, 5-(1,2-Dihydro-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-2thiophenecarbonitrile, 5-(3-Fluoro-4-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(2-Amino-5-pyrimidinyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile, 5-(3-chlorophenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-Benzyl-5-(3-chlorophenyl)-3-methyl-1,3-dihydro-2H-indole-2-thione, 4-(3,3-dimethyl-2-thioxo-2,3-dihydro-1H-indol-5-yl)-2-furonitrile, 5-(3-methoxyphenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-pyridinecarbonitrile, 5-(3,4-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(5-Chloro-2-thienyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-furancarbonitrile, 5-(3-Chloro-4-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chloro-5-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3,5-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-propyl-2-thiophenecarbonitrile, 5-(3-Fluoro-4-nitrophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-furancarbonitrile, 5''-(3-Chlorophenyl)spiro[cyclobutane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(2-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(4-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-4-methyl-2-thiophenecarbonitrile, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-2-thiophenecarbonitrile, 5''-(3-Fluorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-

AHPWA22AUSA

2''(1''H)-thione, 5-(3-Hydroxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-chlorophenyl)-3,3-diethyl-1,3-dihydro-2H-indole-2-thione, 5-(4-Fluoro-3-(trifluoromethyl)phenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-n-butyl-2-thiophenecarbonitrile, 5-(3-Fluoro-5-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chlorophenyl)-N-hydroxyspiro[cyclohexane-1,3'-[3H]indol]-2-amine, N-(Acetyloxy)-5'-(3-chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2''amine, 5'-(3-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(2-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(4-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3,4-difluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-methoxyphenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-nitrophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-cyanophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 3-(1',2'-Dihydro-2'-(hydroxyimino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluorobenzonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-1-methyl-2-carbonitrile, 5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-2-carbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'(acetoxymino)-5'-yl)-2-thiophenecarbonitrile, 3-Fluoro-N'-hydroxy-5-(2'-(hydroxyamino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarboximidamide, N'-Hydroxy-4-(spiro[cyclohexane-1,3'-[3H]indol]-2'-hydroxyimino)-5'-yl-2-thiophenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarboxidamide,

AHPWA22AUSA

5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(3-Cyano-5-fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-3-methyl-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-thiophen-3-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 3-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluoro-benzonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-4-methyl-thiophene-2-carbonitrile, and 4-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

25(Original). The method according to claim 23, wherein said compound is 5'-(5-Cyano-1-methyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

BEST AVAILABLE COPY